

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
27 July 2000 (27.07.2000)

PCT

(10) International Publication Number
WO 00/43373 A3(51) International Patent Classification⁷: C07D 239/47,
239/88, 239/93, 239/95, A61K 31/513, 31/517, A61P
11/00, 25/00, 35/00, 37/00[US/US]; 90 Poor Street, Andover, MA 01810 (US).
TOLEDO-SHERMAN, Leticia, M. [CU/US]; 176 Sum-
mer Street #3, Somerville, MA 02143 (US).

(21) International Application Number: PCT/US00/01581

(74) Agents: TSAO, Y., Rocky et al.; Fish & Richardson P.C.,
225 Franklin Street, Boston, MA 02110-2804 (US).

(22) International Filing Date: 21 January 2000 (21.01.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/116,697 22 January 1999 (22.01.1999) US(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,
UG, US, UZ, VN, YU, ZA, ZW.(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US 60/116,697 (CIP)
Filed on 22 January 1999 (22.01.1999)(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GW, ML, MR, NE, SN, TD, TG).(71) Applicant (*for all designated States except US*):
KINETIX PHARMACEUTICALS, INC. [US/US];
Suite 3500, 200 Boston Avenue, Medford, MA 02155
(US).

Published:

- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): ARMISTEAD,
David, M. [US/US]; 10 Briant Drive, Sudbury, MA 01776
(US). BEMIS, Jean, E. [US/US]; 256 Appleton Street,
Arlington, MA 02476 (US). ELBAUM, Daniel [US/US];
25 Sherrin Road, Newton, MA 02462 (US). HABGOOD,
Gregory [US/US]; 21 Carriage Court, Merrimac, MA
01860 (US). NOVAK, Perry, M. [US/US]; 35 Debbie
Lane, Milford, MA 01757 (US). NUNES, Joseph, J.(88) Date of publication of the international search report:
28 December 2000For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: KINASE INHIBITORS

(57) Abstract: The invention relates to inhibitors of kinases, compositions comprising the inhibitors, and methods of using the inhibitors and inhibitor compositions. The inhibitors and compositions comprising them are useful for treating disease or disease symptoms. The invention also provides for methods of making kinase inhibitor compounds, methods of inhibiting kinase activity, and methods for treating disease or disease symptoms.

WO 00/43373 A3

INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 00/01581

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/47 C07D239/88 C07D239/93 C07D239/95 A61K31/513
A61K31/517 A61P11/00 A61P25/00 A61P35/00 A61P37/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 2 048 250 A (THE UPJOHN CO.) 10 December 1980 (1980-12-10) page 25, line 35 - line 41 page 26, line 26 - line 56 claims ---	1-6,8
X	US 4 308 272 A (W. WIERENGA ET AL.) 29 December 1981 (1981-12-29) column 2, line 21 - line 28; claim 1 ---	1-6,8
X	WO 98 25596 A (PHARMACIA & UPJOHN CO.) 18 June 1998 (1998-06-18) page 8, line 31 -page 99, line 23; claims 1,3,8 ---	1-6,8
X	US 4 689 328 A (I. H. HALL ET AL.) 25 August 1987 (1987-08-25) claims 1-8; table 1 ---	1,2,6,8
-/-		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

26 July 2000

Date of mailing of the international search report

24. 10. 00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hass, C

INTERNATIONAL SEARCH REPORT

Internat. Application No.

PCT/US 00/01581

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PATENT ABSTRACTS OF JAPAN vol. 15, no. 154 (C-0825), 18 April 1991 (1991-04-18) & JP 03 031267 A (MORISHITA PHARMACEUT CO LTD), 12 February 1991 (1991-02-12) abstract ---	1,2,6,8
X	PATENT ABSTRACTS OF JAPAN vol. 11, no. 38 (C-401), 4 February 1987 (1987-02-04) & JP 61 205260 A (ZERIA SHINYAKU KOGYO KK), 11 September 1986 (1986-09-11) abstract; formulae I, II ---	1-6,18, 19
X	V. J. RAM ET AL.: EUROPEAN JOURNAL OF MEDICINAL CHEMISTRY, vol. 25, no. 6, 1990, pages 533-8, XP000929367 page 534, compounds 21 and 3 to 14 ---	1-6,18, 19
X	S. A. ABDEL-AZIZ ET AL.: PHOSPHORUS, SULFUR AND SILICON, vol. 113, no. 1-4, 1996, pages 67-77, XP000864775 page 68, compounds (1) and (4); page 70, compound (10) ---	1-6,18
X	N. RASHED ET AL.: JOURNAL OF THE CHINESE CHEMICAL SOCIETY, vol. 40, no. 4, 1993, pages 393-7, XP000864792 Taipei page 394, scheme I, compounds 1, 5; page 395, compound 5 ---	1-6,18
X	GB 2 107 309 A (EGYT GYOGYSZERVEGYESZETI GYAR) 27 April 1983 (1983-04-27) examples 1-4 ---	1-6
X	GB 2 158 068 A (ASTON MOLECULES LTD.) 6 November 1985 (1985-11-06) page 1, line 19 - line 33 ---	1-6
X	WO 98 14504 A (DSM N. V.) 9 April 1998 (1998-04-09) example V ---	1-6
X	US 3 378 557 A (K. J. RORIG ET AL.) 16 April 1968 (1968-04-16) column 3, line 60 ---	1-6
	--- -/--	

INTERNATIONAL SEARCH REPORT

Intern: 1st Application No

PCT/US 00/01581

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	M. F. M. HASSAN ET AL.: CHINESE JOURNAL OF CHEMISTRY, vol. 9, no. 3, 1991, pages 262-9, XP000925588 page 264, compounds 1, 2, 5, 6, 7 ---	1-6
X	V. J. RAM: JOURNAL FÜR PRAKTISCHE CHEMIE, vol. 331, no. 6, 1989, pages 957-63, XP000864778 page 958, compound 15 ---	1-6
X	V. J. RAM ET AL.: JOURNAL FÜR PRAKTISCHE CHEMIE, vol. 332, no. 5, 1990, pages 629-39, XP000864779 page 631, scheme 2, compounds 14 a-d; page 630, scheme 1, definition of QR ---	1-6
X	V. J. RAM: JOURNAL FÜR PRAKTISCHE CHEMIE, vol. 331, no. 6, 1989, pages 893-905, XP000864780 page 894, compounds 6-17 ---	1-6
X	V. J. RAM: ARCHIV DER PHARMAZIE, vol. 323, no. 11, 1990, pages 895-9, XP000864784 scheme 1 ---	1-6
X	N. H. ESHBA: ALEXANDRIA JOURNAL OF PHARMACEUTICAL SCIENCE, vol. 9, no. 1, 1995, pages 31-4, XP000864773 page 32, scheme 1, first formula, compounds VII-IX ---	1-6
X	F. A. ATTABY ET AL.: ARCH. PHARMACAL RES., vol. 20, no. 6, 1997, pages 620-8, XP000864772 page 625, compounds 1a-c, 2a-c; page 627, compounds 20a-c, 27a-f, 24a-c ---	1-6
X	S. KAMBE ET AL.: SYNTHESIS, no. 4, 1979, pages 287-9, XP000929368 page 288, compounds 4a, 8 ---	1-6
X	S. M. HUSSAIN ET AL.: JOURNAL OF HETEROCYCLIC CHEMISTRY, vol. 24, no. 6, 1987, pages 1605-10, XP000929369 page 1605, compounds I, II; page 1607, compounds XIX, XXI; page 1608, compound XXIII ---	1-6

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/01581

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	S. M. HUSSAIN ET AL.: JOURNAL OF HETEROCYCLIC CHEMISTRY, vol. 22, no. 1, 1985, pages 169-71, XP000929371 page 169, compounds IV, VI; page 170, compound IX ---	1-6
X	A. M. EL-REEDY ET AL.: JOURNAL OF HETEROCYCLIC CHEMISTRY, vol. 26, no. 2, 1989, pages 313-6, XP000929372 page 313, compounds I, III ---	1-6
X	V. J. RAM ET AL.: LIEBIGS ANNALEN DER CHEMIE, no. 9, 1987, pages 797-801, XP000929326 page 798, compounds 1a-1d, 5a-5c ---	1-6
P,X	J. M. PARMAR ET AL.: INDIAN JOURNAL OF CHEMISTRY, vol. 38b, no. 4, April 1999 (1999-04), pages 440-4, XP000864783 page 441, compound 1 ---	1-6
X	J. F. W. KEANA ET AL.: JOURNAL OF ORGANIC CHEMISTRY, vol. 41, no. 12, 1976, pages 2124-9, XP002143538 page 2125, chart 1, compound 3 ---	1,2,6
	M. ENDUKALIA ET AL.: CHEMICAL AND	1,2,6

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 00/01581

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 10-15 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-6, 8(partly), 9(partly), 10-19

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-6, 8(partly), 9(partly), 10-19

Pyrimidine derivatives of the formula as defined in claim 1,
their pharmaceutical use and a process for their preparation.

2. Claims: 7, 8(partly), 9(partly)

Condensed pyrimidine derivatives of the formula as defined
in claim 7 and pharmaceutical compositions comprising them.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/01581

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 2048250	A	10-12-1980	BE 882315 A	19-09-1980
			CH 646958 A	28-12-1984
			DE 3008693 A	02-10-1980
			FR 2451918 A	17-10-1980
			IT 1143012 B	22-10-1986
			JP 1797601 C	28-10-1993
			JP 5002670 B	13-01-1993
			JP 55127378 A	02-10-1980
			JP 1905169 C	08-02-1995
			JP 5017451 A	26-01-1993
			JP 6027070 B	13-04-1994
			NL 8001568 A	23-09-1980
			US 5434157 A	18-07-1995
			US 5002951 A	26-03-1991
			US 4507302 A	26-03-1985
			US 4543248 A	24-09-1985
			US 4665077 A	12-05-1987
			US 5554617 A	10-09-1996
			US 4619933 A	28-10-1986

US 4308272	A	29-12-1981	NONE	

WO 9825596	A	18-06-1998	AU 5687198 A	03-07-1998
			EP 0948331 A	13-10-1999

US 4689328	A	25-08-1987	NONE	

JP 03031267	A	12-02-1991	NONE	

JP 61205260	A	11-09-1986	NONE	

GB 2107309	A	27-04-1983	HU 187355 B	28-12-1985
			BE 894193 A	25-02-1983
			CH 648833 A	15-04-1985
			ES 515190 D	01-06-1983
			FI 822818 A	01-03-1983
			FR 2512021 A	04-03-1983
			IT 1190971 B	24-02-1988
			JP 58074671 A	06-05-1983
			SU 1132789 A	30-12-1984

GB 2158068	A	06-11-1985	NONE	

WO 9814504	A	09-04-1998	NL 1004192 C	07-04-1998
			AU 4401897 A	24-04-1998
			EP 0929597 A	21-07-1999

US 3378557	A	16-04-1968	FR 3765 M	
			GB 1003802 A	

WO 9733883	A	18-09-1997	EP 0888335 A	07-01-1999
			JP 2000506532 T	30-05-2000

WO 9818782	A	07-05-1998	AU 4954097 A	22-05-1998
			EP 0934304 A	11-08-1999
			US 6114333 A	05-09-2000

WO 9744326	A	27-11-1997	AU 2897897 A	09-12-1997

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/01581

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9744326 A		BR 9709599 A	10-08-1999
		CZ 9803803 A	14-04-1999
		EP 0901474 A	17-03-1999
		HR 970275 A	30-04-1998
		HU 9901535 A	28-09-1999
		NO 985392 A	25-01-1999
		PL 330069 A	26-04-1999
		US 5863924 A	26-01-1999
		US 5952331 A	14-09-1999
		US 5958934 A	28-09-1999

C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF₃, OR¹³, SR¹³, NR¹³R¹³, COOR¹³, NO₂, CN, C(O)R¹³, C(O)NR¹³R¹³, NHC(O)R¹³, or OC(O)R¹³; or phenyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF₃, OR¹³, SR¹³, NR¹³R¹³, COOR¹³, NO₂, CN, C(O)R¹³, C(O)NR¹³R¹³, NHC(O)R¹³, or OC(O)R¹³;

Each R¹¹ is independently C(O)R¹⁰, COOR¹⁰, or S(O)₂R¹⁰;

Each R¹² is independently H, C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, or phenyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF₃, OR¹³, SR¹³, NR¹³R¹³, COOR¹³, NO₂, CN, C(O)R¹³, C(O)NR¹³R¹³, NHC(O)R¹³, or OC(O)R¹³;

Each R¹³ is independently H; C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; C1-C10 alkyl optionally substituted with halo, CF₃, OR¹⁹, SR¹⁹, NR¹⁹R¹⁹, COOR¹⁹, NO₂, CN; or phenyl optionally substituted with halo, CF₃, OR¹⁹, SR¹⁹, NR¹⁹R¹⁹, COOR¹⁹, NO₂, CN;


Each R¹⁵ is independently C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R⁹; C1-C10 alkyl substituted with 1-3 independent aryl, R⁷ or R⁹ groups; or C1-C10 alkenyl substituted with 1-3 independent aryl, R⁷ or R⁹;

Each R¹⁶ is independently C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; R⁹; C1-C10 alkyl substituted with 1-3 independent aryl, R⁷ or R⁹ groups; C1-C10 alkenyl substituted with 1-3 independent aryl, R⁷ or R⁹; or phenyl substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, R⁹, halo, CF₃, OR¹², SR¹², NR¹²R¹², COOR¹², NO₂, CN, C(O)R¹², C(O)NR¹²R¹², NHC(O)R¹², NH(COOR¹²), S(O)₂NR¹²R¹², OC(O)R¹², C1-C10 alkyl substituted with 1-3 independent R⁹, halo, CF₃, OR¹², SR¹², NR¹²R¹², COOR¹², NO₂, CN, C(O)R¹², C(O)NR¹²R¹², NHC(O)R¹²

Each haloalkyl is independently a C1-C10 alkyl substituted with one or more halogen atoms, selected from F, Cl, Br, or I, wherein the number of halogen atoms may not exceed that number that results in a perhaloalkyl group; and

Each aryl is independently a 6-carbon monocyclic or 10-carbon bicyclic aromatic ring system optionally substituted with 1-3 independent C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; R^9 ; halo; haloalkyl; CF_3 ; OR^{12} ; SR^{12} ; $NR^{12}R^{12}$; $COOR^{12}$; NO_2 ; CN ; $C(O)R^{12}$; $C(O)C(O)R^{12}$; $C(O)NR^{12}R^{12}$; $S(O)_2R^{12}$; $N(R^{12})C(O)R^{12}$; $N(R^{12})(COOR^{12})$; $N(R^{12})S(O)_2R^{12}$; $S(O)_2NR^{12}R^{12}$; $OC(O)R^{12}$; $NR^{12}C(O)NR^{12}R^{12}$; $NR^{12}C(O)C(O)R^{12}$; $NR^{12}C(O)R^9$; $NR^{12}S(O)_2NR^{12}R^{12}$; $NR^{12}S(O)_2R^9$; $NR^{12}C(O)C(O)NR^{12}R^{12}$; C1-C10 alkyl substituted with 1-3 independent R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN , $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$; C2-C10 alkenyl substituted with 1-3 independent R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN , $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$; or R^{12} .

2. The compound of claim 1 having the formula:



wherein,

R^1 is H; $COOR^5$; $C(O)NR^5R^5$; halo; C2-C10 alkyl; C1-C10 alkenyl; C1-C10 alkyl substituted with NR^5R^5 , NR^5R^6 , SR^5 or OR^5 ; or C1-C10 alkenyl substituted with NR^5R^5 , NR^5R^6 , SR^5 or OR^5 ;

R^2 is NR^5R^{15} ; SR^5 ; OR^5 ; R^8 ; aryl; $N(R^5)-N=CH(R^8)$; $N(R^5)-N=CH(aryl)$; $NR^5-NR^5C(O)NR^5R^5$; $NR^5-NR^5R^{16}$; $NR^5-NR^5R^6$; C1-C10 alkyl substituted with aryl, R^8 , halo, CF_3 , SR^5 , OR^5 , $OC(O)R^5$, NR^5R^5 , NR^5R^6 , $COOR^5$, NO_2 , CN , $C(O)R^5$, $C(O)NR^5R^5$, or $S(O)_2NR^5R^5$; or C1-C10 alkenyl substituted with aryl, R^8 , halo, CF_3 ,

Each R^9 is independently a 5-8 membered monocyclic, 8-12 membered bicyclic, or 11-14 membered tricyclic ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if bicyclic, or 1-9 heteroatoms if tricyclic, said heteroatoms selected from O, N, or S, which may be saturated or unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent independently selected from C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; halo; sulfur; oxygen; CF_3 ; haloalkyl; SR^{10} ; OR^{10} ; $NR^{10}R^{10}$; $NR^{10}R^{11}$; $NR^{11}R^{11}$; $COOR^{10}$; NO_2 ; CN ; $C(O)R^{10}$; or $C(O)NR^{10}R^{10}$;

Each R^{10} is independently H, C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; haloalkyl; C1-C10 alkyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF_3 , OR^{13} , SR^{13} , $NR^{13}R^{13}$, $COOR^{13}$, NO_2 , CN , $C(O)R^{13}$, $C(O)NR^{13}R^{13}$, $NHC(O)R^{13}$, or $OC(O)R^{13}$; or phenyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF_3 , OR^{13} , SR^{13} , $NR^{13}R^{13}$, $COOR^{13}$, NO_2 , CN , $C(O)R^{13}$, $C(O)NR^{13}R^{13}$, $NHC(O)R^{13}$, or $OC(O)R^{13}$;

Each R^{11} is independently $C(O)R^{10}$, $COOR^{10}$, or $S(O)_2R^{10}$;

Each R^{12} is independently H, C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, or phenyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF_3 , OR^{13} , SR^{13} , $NR^{13}R^{13}$, $COOR^{13}$, NO_2 , CN , $C(O)R^{13}$, $C(O)NR^{13}R^{13}$, $NHC(O)R^{13}$, or $OC(O)R^{13}$;

Each R^{13} is independently H; C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; C1-C10 alkyl optionally substituted with halo, CF_3 , OR^{19} , SR^{19} , $NR^{19}R^{19}$, $COOR^{19}$, NO_2 , CN ; or phenyl optionally substituted with halo, CF_3 , OR^{19} , SR^{19} , $NR^{19}R^{19}$, $COOR^{19}$, NO_2 , CN ;


Each R^{15} is independently C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^9 ; C1-C10 alkyl substituted with one or two independent aryl, R^7 or R^9 groups; or C1-C10 alkenyl substituted with aryl, R^7 or R^9 ;

Each R^{16} is independently C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; R^9 ; C1-C10 alkyl substituted with one or two independent aryl, R^7 or R^9 groups; C1-C10 alkenyl substituted with aryl, R^7 or R^9 ; or phenyl substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN , $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$, C1-C10 alkyl substituted with R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN , $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$;

Each R^{19} is independently H; C1-C10 alkyl; C3-C10 cycloalkyl or phenyl;

Each haloalkyl is independently a C1-C10 alkyl substituted with one or more halogen atoms, selected from F, Cl, Br, or I, wherein the number of halogen atoms may not exceed that number that results in a perhaloalkyl group;

Each aryl is independently a 6-carbon monocyclic or 10-carbon bicyclic aromatic ring system optionally substituted with 1-3 independent C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; R^9 ; halo; haloalkyl; CF_3 ; OR^{12} ; SR^{12} ; $NR^{12}R^{12}$; $COOR^{12}$; NO_2 ; CN ; $C(O)R^{12}$; $C(O)C(O)R^{12}$; $C(O)NR^{12}R^{12}$; $S(O)_2R^{12}$; $N(R^{12})C(O)R^{12}$; $N(R^{12})(COOR^{12})$; $N(R^{12})S(O)_2R^{12}$; $S(O)_2NR^{12}R^{12}$; $OC(O)R^{12}$; $NR^{12}C(O)NR^{12}R^{12}$; $NR^{12}C(O)C(O)R^{12}$; $NR^{12}C(O)R^9$; $NR^{12}S(O)_2NR^{12}R^{12}$; $NR^{12}S(O)_2R^9$; $NR^{12}C(O)C(O)NR^{12}R^{12}$; C1-C10 alkyl substituted with 1-3 independent R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN , $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$; C2-C10 alkenyl substituted with



wherein,

R^1 is CN;

R^2 is NR^5R^{15} ; OR^5 ; R^8 ; aryl; $N(R^5)-N=CH(R^8)$; $N(R^5)-N=CH(aryl)$; $NR^5-NR^5C(O)NR^5R^5$; $NR^5-NR^5R^{16}$; $NR^5-NR^5R^6$; C1-C10 alkyl substituted with aryl, R^8 , halo, CF_3 , SR^5 , OR^5 , $OC(O)R^5$, NR^5R^5 , NR^5R^6 , $COOR^5$, NO_2 , CN , $C(O)R^5$, $C(O)NR^5R^5$, or $S(O)_2NR^5R^5$; or C1-C10 alkenyl substituted with aryl, R^8 , halo, CF_3 , SR^5 , OR^5 , $OC(O)R^5$, NR^5R^5 , NR^5R^6 , $COOR^5$, NO_2 , CN , $C(O)R^5$, $C(O)NR^5R^5$, or $S(O)_2NR^5R^5$;

X is O or S;

Each R^4 is independently selected from H, C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^8 ; halo; haloalkyl; CF_3 ; SR^5 ; OR^5 ; NR^5R^5 ; NR^5R^6 ; $COOR^5$; NO_2 ; CN ; $C(O)R^5$; $C(O)C(O)R^5$; $C(O)NR^5R^5$; $OC(O)R^5$; $S(O)_2R^5$; $S(O)_2NR^5R^5$; $NR^5C(O)NR^5R^5$; $NR^5C(O)C(O)R^5$; $NR^5C(O)R^5$; $NR^5(COOR^5)$; $NR^5C(O)R^8$; $NR^5S(O)_2NR^5R^5$; $NR^5S(O)_2R^5$; $NR^5S(O)_2R^8$; $NR^5C(O)C(O)NR^5R^5$; $NR^5C(O)C(O)NR^5R^6$; C1-C10 alkyl substituted with aryl, R^7 or R^8 ; or C1-C10 alkenyl substituted with aryl, R^7 or R^8 ;

Each R^5 is independently H; C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^9 ; C1-C10 alkyl substituted with one or two independent aryl, R^7 or R^9 groups; C3-C10 cycloalkyl substituted with one or two independent aryl, R^7 or R^9 groups; or C1-C10 alkenyl substituted with aryl, R^7 or R^9 ;

Each R^6 is independently $C(O)R^5$, $COOR^5$, or $S(O)_2R^5$;

Each R^7 is independently halo, CF_3 , SR^{10} , OR^{10} , $OC(O)R^{10}$, $NR^{10}R^{10}$, $NR^{10}R^{11}$, $NR^{11}R^{11}$, $COOR^{10}$, NO_2 , CN , $C(O)R^{10}$, or $C(O)NR^{10}R^{10}$;

Each R^8 is independently a 5-8 membered monocyclic, 8-12 membered bicyclic, or 11-14 membered tricyclic ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if bicyclic, or 1-9 heteroatoms if tricyclic, said heteroatoms selected from O, N, or S, which may be saturated or unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent independently selected from C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^9 ; halo; sulfur; oxygen; CF_3 ; haloalkyl; SR^5 ; OR^5 ; $OC(O)R^5$; NR^5R^5 ; NR^5R^6 ; NR^6R^6 ; $COOR^5$; NO_2 ; CN ; $C(O)R^5$; $C(O)NR^5R^5$; C1-C10 alkyl substituted with R^7 , R^9 or aryl; C1-C10 alkenyl substituted with R^7 , R^9 or aryl;

Each R^9 is independently a 5-8 membered monocyclic, 8-12 membered bicyclic, or 11-14 membered tricyclic ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if bicyclic, or 1-9 heteroatoms if tricyclic, said heteroatoms selected from O, N, or S, which may be saturated or unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent independently selected from C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; halo; sulfur; oxygen; CF_3 ; haloalkyl; SR^{10} ; OR^{10} ; $NR^{10}R^{10}$; $NR^{10}R^{11}$; $NR^{11}R^{11}$; $COOR^{10}$; NO_2 ; CN ; $C(O)R^{10}$; or $C(O)NR^{10}R^{10}$;

Each R^{10} is independently H, C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; haloalkyl; C1-C10 alkyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF_3 , OR^{13} , SR^{13} , $NR^{13}R^{13}$, $COOR^{13}$, NO_2 , CN , $C(O)R^{13}$, $C(O)NR^{13}R^{13}$, $NHC(O)R^{13}$, or $OC(O)R^{13}$; or phenyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF_3 , OR^{13} , SR^{13} , $NR^{13}R^{13}$, $COOR^{13}$, NO_2 , CN , $C(O)R^{13}$, $C(O)NR^{13}R^{13}$, $NHC(O)R^{13}$, or $OC(O)R^{13}$;

Each R^{11} is independently $C(O)R^{10}$, $COOR^{10}$, or $S(O)_2R^{10}$;

Each R^{12} is independently H, C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF_3 , OR^{13} , SR^{13} , $NR^{13}R^{13}$, $COOR^{13}$, NO_2 , CN , $C(O)R^{13}$, $C(O)NR^{13}R^{13}$, $NHC(O)R^{13}$, or $OC(O)R^{13}$;

C10 cycloalkenyl, halo, CF₃, OR¹³, SR¹³, NR¹³R¹³, COOR¹³, NO₂, CN, C(O)R¹³, C(O)NR¹³R¹³, NHC(O)R¹³, or OC(O)R¹³;

Each R¹³ is independently H; C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; C1-C10 alkyl optionally substituted with halo, CF₃, OR¹⁹, SR¹⁹, NR¹⁹R¹⁹, COOR¹⁹, NO₂, CN; or phenyl optionally substituted with halo, CF₃, OR¹⁹, SR¹⁹, NR¹⁹R¹⁹, COOR¹⁹, NO₂, CN;

Each R¹⁴ is each independently selected from C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R⁸; halo; haloalkyl; CF₃; SR⁵; OR⁵; NR⁵R⁵; NR⁵R⁶; COOR⁵; NO₂; CN; C(O)R⁵; C(O)C(O)R⁵; C(O)NR⁵R⁵; OC(O)R⁵; S(O)₂R⁵; S(O)₂NR⁵R⁵; NR⁵C(O)NR⁵R⁵; NR⁵C(O)C(O)R⁵; NR⁵C(O)R⁵; NR⁵(COOR⁵); NR⁵C(O)R⁸; NR⁵S(O)₂NR⁵R⁵; NR⁵S(O)₂R⁵; NR⁵S(O)₂R⁸; NR⁵C(O)C(O)NR⁵R⁵; NR⁵C(O)C(O)NR⁵R⁶; C1-C10 alkyl substituted with aryl, R⁷ or R⁸; or C1-C10 alkenyl substituted with aryl, R⁷ or R⁸;

Each R¹⁵ is independently C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R⁹; C1-C10 alkyl substituted with one or two independent aryl, R⁷ or R⁹ groups; or C1-C10 alkenyl substituted with aryl, R⁷ or R⁹;

Each R¹⁶ is independently C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; R⁹; C1-C10 alkyl substituted with one or two independent aryl, R⁷ or R⁹ groups; C1-C10 alkenyl substituted with aryl, R⁷ or R⁹; or phenyl substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, R⁹, halo, CF₃, OR¹², SR¹², NR¹²R¹², COOR¹², NO₂, CN, C(O)R¹², C(O)NR¹²R¹², NHC(O)R¹², NH(COOR¹²), S(O)₂NR¹²R¹², OC(O)R¹², C1-C10 alkyl substituted with R⁹, halo, CF₃, OR¹², SR¹², NR¹²R¹², COOR¹², NO₂, CN, C(O)R¹², C(O)NR¹²R¹², NHC(O)R¹², NH(COOR¹²), S(O)₂NR¹²R¹², OC(O)R¹²;

Each R¹⁷ is independently selected from H, C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R⁸; halo; haloalkyl; CF₃; SR⁵; OR⁵; NR⁵R⁵; NR⁵R⁶; COOR⁵; NO₂; CN; C(O)R⁵; C(O)C(O)R⁵;

$C(O)NR^5R^5$; $OC(O)R^5$; $S(O)_2R^5$; $S(O)_2NR^5R^5$; $NR^5C(O)NR^5R^5$; $NR^5C(O)C(O)R^5$; $NR^5C(O)R^5$; $NR^5(COOR^5)$; $NR^5C(O)R^8$; $NR^5S(O)_2NR^5R^5$; $NR^5S(O)_2R^5$; $NR^5S(O)_2R^8$; $NR^5C(O)C(O)NR^5R^5$; $NR^5C(O)C(O)NR^5R^6$; C1-C10 alkyl substituted with aryl, R^7 or R^8 ; or C1-C10 alkenyl substituted with aryl, R^7 or R^8 ;

Each R^{19} is independently H; C1-C10 alkyl; C3-C10 cycloalkyl or phenyl;

Each haloalkyl is independently a C1-C10 alkyl substituted with one or more halogen atoms, selected from F, Cl, Br, or I, wherein the number of halogen atoms may not exceed that number that results in a perhaloalkyl group;

Each aryl is independently a 6-carbon monocyclic or 10-carbon bicyclic aromatic ring system optionally substituted with 1-3 independent C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; R^9 ; halo; haloalkyl; CF_3 ; OR^{12} ; SR^{12} ; $NR^{12}R^{12}$; $COOR^{12}$; NO_2 ; CN ; $C(O)R^{12}$; $C(O)C(O)R^{12}$; $C(O)NR^{12}R^{12}$; $S(O)_2R^{12}$; $N(R^{12})C(O)R^{12}$; $N(R^{12})(COOR^{12})$; $N(R^{12})S(O)_2R^{12}$; $S(O)_2NR^{12}R^{12}$; $OC(O)R^{12}$; $NR^{12}C(O)NR^{12}R^{12}$; $NR^{12}C(O)C(O)R^{12}$; $NR^{12}C(O)R^9$; $NR^{12}S(O)_2NR^{12}R^{12}$; $NR^{12}S(O)_2R^9$; $NR^{12}C(O)C(O)NR^{12}R^{12}$; C1-C10 alkyl substituted with 1-3 independent R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN , $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$; C2-C10 alkenyl substituted with 1-3 independent R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN , $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$; or R^{12} ;

wherein when all R^4 and R^{17} are simultaneously H, R^{14} may not be Me, Cl, or OMe; and wherein R^{14} and R^{17} may not simultaneously be Cl.

4. The compound of claim 1 having the formula,

wherein,

R^1 is CN;

R^2 is NR^5R^{15} ; OR^5 ; R^8 ; aryl; $N(R^5)-N=CH(R^8)$; $N(R^5)-N=CH(aryl)$; $NR^5-NR^5C(O)NR^5R^5$; $NR^5-NR^5R^{16}$; $NR^5-NR^5R^6$; C1-C10 alkyl substituted with aryl, R^8 , halo, CF_3 , SR^5 , OR^5 , $OC(O)R^5$, NR^5R^5 , NR^5R^6 , $COOR^5$, NO_2 , CN, $C(O)R^5$, $C(O)NR^5R^5$, or $S(O)_2NR^5R^5$; or C1-C10 alkenyl substituted with aryl, R^8 , halo, CF_3 , SR^5 , OR^5 , $OC(O)R^5$, NR^5R^5 , NR^5R^6 , $COOR^5$, NO_2 , CN, $C(O)R^5$, $C(O)NR^5R^5$, or $S(O)_2NR^5R^5$;

X is O or S;

Each R^4 is independently selected from H, C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^8 ; halo; haloalkyl; CF_3 ; SR^5 ; OR^5 ; NR^5R^5 ; NR^5R^6 ; $COOR^5$; NO_2 ; CN; $C(O)R^5$; $C(O)C(O)R^5$; $C(O)NR^5R^5$; $OC(O)R^5$; $S(O)_2R^5$; $S(O)_2NR^5R^5$; $NR^5C(O)NR^5R^5$; $NR^5C(O)C(O)R^5$; $NR^5C(O)R^5$; $NR^5(COOR^5)$; $NR^5C(O)R^8$; $NR^5S(O)_2NR^5R^5$; $NR^5S(O)_2R^5$; $NR^5S(O)_2R^8$; $NR^5C(O)C(O)NR^5R^5$; $NR^5C(O)C(O)NR^5R^6$; C1-C10 alkyl substituted with aryl, R^7 or R^8 ; or C1-C10 alkenyl substituted with aryl, R^7 or R^8 ;

Each R^5 is independently H; C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^9 ; C1-C10 alkyl substituted with one or two independent aryl, R^7 or R^9 groups; C3-C10 cycloalkyl substituted with one or two independent aryl, R^7 or R^9 groups; or C1-C10 alkenyl substituted with aryl, R^7 or R^9 ;

Each R^6 is independently $C(O)R^5$, $COOR^5$, or $S(O)_2R^5$;

Each R^7 is independently halo, CF_3 , SR^{10} , OR^{10} , $OC(O)R^{10}$, $NR^{10}R^{10}$, $NR^{10}R^{11}$, $NR^{11}R^{11}$, $COOR^{10}$, NO_2 , CN, $C(O)R^{10}$, or $C(O)NR^{10}R^{10}$;

Each R^8 is independently a 5-8 membered monocyclic, 8-12 membered bicyclic, or 11-14 membered tricyclic ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if bicyclic, or 1-9 heteroatoms if tricyclic, said heteroatoms selected from O, N, or S, which may be saturated or unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent independently selected from C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl;

aryl; R^9 ; halo; sulfur; oxygen; CF_3 ; haloalkyl; SR^5 ; OR^5 ; $OC(O)R^5$; NR^5R^5 ; NR^5R^6 ; NR^6R^6 ; $COOR^5$; NO_2 ; CN ; $C(O)R^5$; $C(O)NR^5R^5$; C1-C10 alkyl substituted with R^7 , R^9 or aryl; C1-C10 alkenyl substituted with R^7 , R^9 or aryl;

Each R^9 is independently a 5-8 membered monocyclic, 8-12 membered bicyclic, or 11-14 membered tricyclic ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if bicyclic, or 1-9 heteroatoms if tricyclic, said heteroatoms selected from O, N, or S, which may be saturated or unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent independently selected from C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; halo; sulfur; oxygen; CF_3 ; haloalkyl; SR^{10} ; OR^{10} ; $NR^{10}R^{10}$; $NR^{10}R^{11}$; $NR^{11}R^{11}$; $COOR^{10}$; NO_2 ; CN ; $C(O)R^{10}$; or $C(O)NR^{10}R^{10}$;

Each R^{10} is independently H, C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; haloalkyl; C1-C10 alkyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF_3 , OR^{13} , SR^{13} , $NR^{13}R^{13}$, $COOR^{13}$, NO_2 , CN , $C(O)R^{13}$, $C(O)NR^{13}R^{13}$, $NHC(O)R^{13}$, or $OC(O)R^{13}$; or phenyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF_3 , OR^{13} , SR^{13} , $NR^{13}R^{13}$, $COOR^{13}$, NO_2 , CN , $C(O)R^{13}$, $C(O)NR^{13}R^{13}$, $NHC(O)R^{13}$, or $OC(O)R^{13}$;

Each R^{11} is independently $C(O)R^{10}$, $COOR^{10}$, or $S(O)_2R^{10}$;

Each R^{12} is independently H, C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, or phenyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF_3 , OR^{13} , SR^{13} , $NR^{13}R^{13}$, $COOR^{13}$, NO_2 , CN , $C(O)R^{13}$

Each R^{14} is independently selected from C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^8 ; halo; haloalkyl; CF_3 ; SR^5 ; OR^5 ; NR^5R^5 ; NR^5R^6 ; $COOR^5$; NO_2 ; CN ; $C(O)R^5$; $C(O)C(O)R^5$; $C(O)NR^5R^5$; $OC(O)R^5$; $S(O)_2R^5$; $S(O)_2NR^5R^5$; $NR^5C(O)NR^5R^5$; $NR^5C(O)C(O)R^5$; $NR^5C(O)R^5$; $NR^5(COOR^5)$; $NR^5C(O)R^8$; $NR^5S(O)_2NR^5R^5$; $NR^5S(O)_2R^5$; $NR^5S(O)_2R^8$; $NR^5C(O)C(O)NR^5R^5$; $NR^5C(O)C(O)NR^5R^6$; C1-C10 alkyl substituted with aryl, R^7 or R^8 ; or C1-C10 alkenyl substituted with aryl, R^7 or R^8 ;

Each R^{15} is independently C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^9 ; C1-C10 alkyl substituted with one or two independent aryl, R^7 or R^9 groups; or C1-C10 alkenyl substituted with aryl, R^7 or R^9 ;

Each R^{16} is independently C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; R^9 ; C1-C10 alkyl substituted with one or two independent aryl, R^7 or R^9 groups; C1-C10 alkenyl substituted with aryl, R^7 or R^9 ; or phenyl substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN , $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$, C1-C10 alkyl substituted with R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN , $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$;


Each R^{19} is independently H; C1-C10 alkyl; C3-C10 cycloalkyl or phenyl;

Each haloalkyl is independently a C1-C10 alkyl substituted with one or more halogen atoms, selected from F, Cl, Br, or I, wherein the number of halogen atoms may not exceed that number that results in a perhaloalkyl group;

Each aryl is independently a 6-carbon monocyclic or 10-carbon bicyclic aromatic ring system optionally substituted with 1-3 independent C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; R^9 ; halo; haloalkyl; CF_3 ; OR^{12} ; SR^{12} ; $NR^{12}R^{12}$; $COOR^{12}$; NO_2 ; CN ; $C(O)R^{12}$; $C(O)C(O)R^{12}$; $C(O)NR^{12}R^{12}$; $S(O)_2R^{12}$; $N(R^{12})C(O)R^{12}$; $N(R^{12})(COOR^{12})$; $N(R^{12})S(O)_2R^{12}$; $S(O)_2NR^{12}R^{12}$; $OC(O)R^{12}$; $NR^{12}C(O)NR^{12}R^{12}$; $NR^{12}C(O)C(O)R^{12}$; $NR^{12}C(O)R^9$; $NR^{12}S(O)_2NR^{12}R^{12}$;

$\text{NR}^{12}\text{S(O)}_2\text{R}^9$; $\text{NR}^{12}\text{C(O)}\text{C(O)}\text{NR}^{12}\text{R}^{12}$; C1-C10 alkyl substituted with 1-3 independent R^9 , halo, CF_3 , OR^{12} , SR^{12} , $\text{NR}^{12}\text{R}^{12}$, COOR^{12} , NO_2 , CN , $\text{C(O)}\text{R}^{12}$, $\text{C(O)}\text{NR}^{12}\text{R}^{12}$, $\text{NHC(O)}\text{R}^{12}$, $\text{NH(COOR}^{12})$, $\text{S(O)}_2\text{NR}^{12}\text{R}^{12}$, $\text{OC(O)}\text{R}^{12}$; C2-C10 alkenyl substituted with 1-3 independent R^9 , halo, CF_3 , OR^{12} , SR^{12} , $\text{NR}^{12}\text{R}^{12}$, COOR^{12} , NO_2 , CN , $\text{C(O)}\text{R}^{12}$, $\text{C(O)}\text{NR}^{12}\text{R}^{12}$, $\text{NHC(O)}\text{R}^{12}$, $\text{NH(COOR}^{12})$, $\text{S(O)}_2\text{NR}^{12}\text{R}^{12}$, $\text{OC(O)}\text{R}^{12}$; or R^{12} ; wherein when all R^4 are H, R^{14} may not be Me or OMe.

5. The compound of claim 1 having the formula,



wherein,

R^1 is CN;

R^2 is NR^5R^{15} ; OR^5 ; R^8 ; aryl; $\text{N(R}^5)\text{-N=CH(R}^8)$; $\text{N(R}^5)\text{-N=CH(aryl)}$; $\text{NR}^5\text{-NR}^5\text{C(O)}\text{NR}^5\text{R}^5$; $\text{NR}^5\text{-NR}^5\text{R}^{16}$; $\text{NR}^5\text{-NR}^5\text{R}^6$; C1-C10 alkyl substituted with aryl, R^8 , halo, CF_3 , SR^5 , OR^5 , $\text{OC(O)}\text{R}^5$, NR^5R^5 , NR^5R^6 , COOR^5 , NO_2 , CN , $\text{C(O)}\text{R}^5$, $\text{C(O)}\text{NR}^5\text{R}^5$, or $\text{S(O)}_2\text{NR}^5\text{R}^5$; or C1-C10 alkenyl substituted with aryl, R^8 , halo, CF_3 , SR^5 , OR^5 , $\text{OC(O)}\text{R}^5$, NR^5R^5 , NR^5R^6 , COOR^5 , NO_2 , CN , $\text{C(O)}\text{R}^5$, $\text{C(O)}\text{NR}^5\text{R}^5$, or $\text{S(O)}_2\text{NR}^5\text{R}^5$;

X is O or S;

Each R^4 is independently selected from H, C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^8 ; halo; haloalkyl; CF_3 ; SR^5 ; OR^5 ; NR^5R^5 ; NR^5R^6 ; COOR^5 ; NO_2 ; CN ; $\text{C(O)}\text{R}^5$; $\text{C(O)}\text{C(O)}\text{R}^5$; $\text{C(O)}\text{NR}^5\text{R}^5$;

$\text{NR}^5\text{C}(\text{O})\text{C}(\text{O})\text{NR}^5\text{R}^5$; $\text{NR}^5\text{C}(\text{O})\text{C}(\text{O})\text{NR}^5\text{R}^6$; C1-C10 alkyl substituted with aryl, R^7 or R^8 ; or C1-C10 alkenyl substituted with aryl, R^7 or R^8 ;

Each R^5 is independently H; C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^9 ; C1-C10 alkyl substituted with one or two independent aryl, R^7 or R^9 groups; C3-C10 cycloalkyl substituted with one or two independent aryl, R^7 or R^9 groups; or C1-C10 alkenyl substituted with aryl, R^7 or R^9 ;

Each R^6 is independently $\text{C}(\text{O})\text{R}^5$, COOR^5 , or $\text{S}(\text{O})_2\text{R}^5$;

Each R^7 is independently halo, CF_3 , SR^{10} , OR^{10} , $\text{OC}(\text{O})\text{R}^{10}$, $\text{NR}^{10}\text{R}^{10}$, $\text{NR}^{10}\text{R}^{11}$, $\text{NR}^{11}\text{R}^{11}$, COOR^{10} , NO_2 , CN , $\text{C}(\text{O})\text{R}^{10}$, or $\text{C}(\text{O})\text{NR}^{10}\text{R}^{10}$;

Each R^8 is independently a 5-8 membered monocyclic, 8-12 membered bicyclic, or 11-14 membered tricyclic ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if bicyclic, or 1-9 heteroatoms if tricyclic, said heteroatoms selected from O, N, or S, which may be saturated or unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent independently selected from C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^9 ; halo; sulfur; oxygen; CF_3 ; haloalkyl; SR^5 ; OR^5 ; $\text{OC}(\text{O})\text{R}^5$; NR^5R^5 ; NR^5R^6 ; NR^6R^6 ; COOR^5 ; NO_2 ; CN ; $\text{C}(\text{O})\text{R}^5$; $\text{C}(\text{O})\text{NR}^5\text{R}^5$; C1-C10 alkyl substituted with R^7 , R^9 or aryl; C1-C10 alkenyl substituted with R^7 , R^9 or aryl;

Each R^9 is independently a 5-8 membered monocyclic, 8-12 membered bicyclic, or 11-14 membered tricyclic ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if bicyclic, or 1-9 heteroatoms if tricyclic, said heteroatoms selected from O, N, or S, which may be saturated or unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent independently selected from C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; halo; sulfur; oxygen; CF_3 ; haloalkyl; SR^{10} ; OR^{10} ; $\text{NR}^{10}\text{R}^{10}$; $\text{NR}^{10}\text{R}^{11}$; $\text{NR}^{11}\text{R}^{11}$; COOR^{10} ; NO_2 ; CN ; $\text{C}(\text{O})\text{R}^{10}$; or $\text{C}(\text{O})\text{NR}^{10}\text{R}^{10}$;

Each R^{10} is independently H, C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; haloalkyl; C1-C10 alkyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-

C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF₃, OR¹³, SR¹³, NR¹³R¹³, COOR¹³, NO₂, CN, C(O)R¹³, C(O)NR¹³R¹³, NHC(O)R¹³, or OC(O)R¹³; or phenyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF₃, OR¹³, SR¹³, NR¹³R¹³, COOR¹³, NO₂, CN, C(O)R¹³, C(O)NR¹³R¹³, NHC(O)R¹³, or OC(O)R¹³;

Each R¹¹ is independently C(O)R¹⁰, COOR¹⁰, or S(O)₂R¹⁰;

Each R¹² is independently H, C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, or phenyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF₃, OR¹³, SR¹³, NR¹³R¹³, COOR¹³, NO₂, CN, C(O)R¹³, C(O)NR¹³R¹³, NHC(O)R¹³, or OC(O)R¹³;

Each R¹³ is independently H; C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; C1-C10 alkyl optionally substituted with halo, CF₃, OR¹⁹, SR¹⁹, NR¹⁹R¹⁹, COOR¹⁹, NO₂, CN; or phenyl optionally substituted with halo, CF₃, OR¹⁹, SR¹⁹, NR¹⁹R¹⁹, COOR¹⁹, NO₂, CN;

Each R¹⁴ is independently selected from C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R⁸; halo; haloalkyl; CF₃; SR⁵; OR⁵; NR⁵R⁵; NR⁵R⁶; COOR⁵; NO₂; CN; C(O)R⁵; C(O)C(O)R⁵; C(O)NR⁵R⁵; OC(O)R⁵; S(O)₂R⁵; S(O)₂NR⁵R⁵; NR⁵C(O)NR⁵R⁵; NR⁵C(O)C(O)R⁵; NR⁵C(O)R⁵; NR⁵(COOR⁵); NR⁵C(O)R⁸; NR⁵S(O)₂NR⁵R⁵; NR⁵S(O)₂R⁵; NR⁵S(O)₂R⁸; NR⁵C(O)C(O)NR⁵R⁵; NR⁵C(O)C(O)NR⁵R⁶; C1-C10 alkyl substituted with aryl, R⁷ or R⁸; or C1-C10 alkenyl substituted with aryl, R⁷ or R⁸;

Each R¹⁵ is independently C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R⁹; C1-C10 alkyl substituted

with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF₃, OR¹³, SR¹³, NR¹³R¹³, COOR¹³, NO₂, CN, C(O)R¹³, C(O)NR¹³R¹³, NHC(O)R¹³, or OC(O)R¹³;

C1-C10 alkyl substituted with

with one or two independent aryl, R^7 or R^9 groups; or C1-C10 alkenyl substituted with aryl, R^7 or R^9 ;

Each R^{17} is independently selected from H, C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^8 ; halo; haloalkyl; CF_3 ; SR^5 ; OR^5 ; NR^5R^5 ; NR^5R^6 ; $COOR^5$; NO_2 ; CN ; $C(O)R^5$; $C(O)C(O)R^5$; $C(O)NR^5R^5$; $OC(O)R^5$; $S(O)_2R^5$; $S(O)_2NR^5R^5$; $NR^5C(O)NR^5R^5$; $NR^5C(O)C(O)R^5$; $NR^5C(O)R^5$; $NR^5(COOR^5)$; $NR^5C(O)R^8$; $NR^5S(O)_2NR^5R^5$; $NR^5S(O)_2R^5$; $NR^5S(O)_2R^8$; $NR^5C(O)C(O)NR^5R^5$; $NR^5C(O)C(O)NR^5R^6$; C1-C10 alkyl substituted with aryl, R^7 or R^8 ; or C1-C10 alkenyl substituted with aryl, R^7 or R^8 ;


Each R^{19} is independently H; C1-C10 alkyl; C3-C10 cycloalkyl or phenyl;

Each haloalkyl is independently a C1-C10 alkyl substituted with one or more halogen atoms, selected from F, Cl, Br, or I, wherein the number of halogen atoms may not exceed that number that results in a perhaloalkyl group;

Each aryl is independently a 6-carbon monocyclic or 10-carbon bicyclic aromatic ring system optionally substituted with 1-3 independent C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; R^9 ; halo; haloalkyl; CF_3 ; OR^{12} ; SR^{12} ; $NR^{12}R^{12}$; $COOR^{12}$; NO_2 ; CN ; $C(O)R^{12}$; $C(O)C(O)R^{12}$; $C(O)NR^{12}R^{12}$; $S(O)_2R^{12}$; $N(R^{12})C(O)R^{12}$; $N(R^{12})(COOR^{12})$; $N(R^{12})S(O)_2R^{12}$; $S(O)_2NR^{12}R^{12}$; $OC(O)R^{12}$; $NR^{12}C(O)NR^{12}R^{12}$; $NR^{12}C(O)C(O)R^{12}$; $NR^{12}C(O)R^9$; $NR^{12}S(O)_2NR^{12}R^{12}$; $NR^{12}S(O)_2R^9$; $NR^{12}C(O)C(O)NR^{12}R^{12}$; C1-C10 alkyl substituted with 1-3 independent R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN , $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$; C2-C10 alkenyl substituted with 1-3 independent R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN , $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$; or R^{12} ;

wherein R^{14} and R^{17} may not simultaneously be Cl.

6. The compound of claim 1 having the formula,



wherein,

R^1 is CN;

R^2 is NR^5R^{15} ; OR^5 ; R^8 ; aryl; $N(R^5)-N=CH(R^8)$; $N(R^5)-N=CH(aryl)$; $NR^5-NR^5C(O)NR^5R^5$; $NR^5-NR^5R^{16}$; $NR^5-NR^5R^6$; C1-C10 alkyl substituted with aryl, R^8 , halo, CF_3 , SR^5 , OR^5 , $OC(O)R^5$, NR^5R^5 , NR^5R^6 , $COOR^5$, NO_2 , CN, $C(O)R^5$, $C(O)NR^5R^5$, or $S(O)_2NR^5R^5$; or C1-C10 alkenyl substituted with aryl, R^8 , halo, CF_3 , SR^5 , OR^5 , $OC(O)R^5$, NR^5R^5 , NR^5R^6 , $COOR^5$, NO_2 , CN, $C(O)R^5$, $C(O)NR^5R^5$, or $S(O)_2NR^5R^5$;

R^3 is R^8 ; $COOR^5$; or C1-C10 alkyl substituted with R^7 , R^8 , or phenyl substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN, $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$, C1-C10 alkyl substituted with R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN, $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$; wherein R^3 is not unsubstituted furanyl, unsubstituted thienyl or unsubstituted pyridyl;

X is O or S;

Each R^4 is independently selected from H, C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^8 ; halo; haloalkyl; CF_3 ; SR^5 ; OR^5 ; NR^5R^5 ; NR^5R^6 ; $COOR^5$; NO_2 ; CN; $C(O)R^5$; $C(O)C(O)R^5$; $C(O)NR^5R^5$; $OC(O)R^5$; $S(O)_2R^5$; $S(O)_2NR^5R^5$; $NR^5C(O)NR^5R^5$; $NR^5C(O)C(O)R^5$; $NR^5C(O)R^5$; $NR^5(COOR^5)$; $NR^5C(O)R^8$; $NR^5S(O)_2NR^5R^5$; $NR^5S(O)_2R^5$; $NR^5S(O)_2R^8$.

$\text{NR}^5\text{C}(\text{O})\text{C}(\text{O})\text{NR}^5\text{R}^5$; $\text{NR}^5\text{C}(\text{O})\text{C}(\text{O})\text{NR}^5\text{R}^6$; C1-C10 alkyl substituted with aryl, R^7 or R^8 ; or C1-C10 alkenyl substituted with aryl, R^7 or R^8 ;

Each R^5 is independently H; C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^9 ; C1-C10 alkyl substituted with one or two independent aryl, R^7 or R^9 groups; C3-C10 cycloalkyl substituted with one or two independent aryl, R^7 or R^9 groups; or C1-C10 alkenyl substituted with aryl, R^7 or R^9 ;

Each R^6 is independently $\text{C}(\text{O})\text{R}^5$, COOR^5 , or $\text{S}(\text{O})_2\text{R}^5$;

Each R^7 is independently halo, CF_3 , SR^{10} , OR^{10} , $\text{OC}(\text{O})\text{R}^{10}$, $\text{NR}^{10}\text{R}^{10}$, $\text{NR}^{10}\text{R}^{11}$, $\text{NR}^{11}\text{R}^{11}$, COOR^{10} , NO_2 , CN , $\text{C}(\text{O})\text{R}^{10}$, or $\text{C}(\text{O})\text{NR}^{10}\text{R}^{10}$;

Each R^8 is independently a 5-8 membered monocyclic, 8-12 membered bicyclic, or 11-14 membered tricyclic ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if bicyclic, or 1-9 heteroatoms if tricyclic, said heteroatoms selected from O, N, or S, which may be saturated or unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent independently selected from C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^9 ; halo; sulfur; oxygen; CF_3 ; haloalkyl; SR^5 ; OR^5 ; $\text{OC}(\text{O})\text{R}^5$; NR^5R^5 ; NR^5R^6 ; NR^6R^6 ; COOR^5 ; NO_2 ; CN ; $\text{C}(\text{O})\text{R}^5$; $\text{C}(\text{O})\text{NR}^5\text{R}^5$; C1-C10 alkyl substituted with R^7 , R^9 or aryl; C1-C10 alkenyl substituted with R^7 , R^9 or aryl;

Each R^9 is independently a 5-8 membered monocyclic, 8-12 membered bicyclic, or 11-14 membered tricyclic ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if bicyclic, or 1-9 heteroatoms if tricyclic, said heteroatoms selected from O, N, or S, which may be saturated or unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent independently selected from C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; halo; sulfur; oxygen; CF_3 ; haloalkyl; SR^{10} ; OR^{10} ; $\text{NR}^{10}\text{R}^{10}$; $\text{NR}^{10}\text{R}^{11}$; $\text{NR}^{11}\text{R}^{11}$; COOR^{10} ; NO_2 ; CN ; $\text{C}(\text{O})\text{R}^{10}$; or $\text{C}(\text{O})\text{NR}^{10}\text{R}^{10}$;

Each R^{10} is independently H, C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; haloalkyl; C1-C10 alkyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-

C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF₃, OR¹³, SR¹³, NR¹³R¹³, COOR¹³, NO₂, CN, C(O)R¹³, C(O)NR¹³R¹³, NHC(O)R¹³, or OC(O)R¹³; or phenyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF₃, OR¹³, SR¹³, NR¹³R¹³, COOR¹³, NO₂, CN, C(O)R¹³, C(O)NR¹³R¹³, NHC(O)R¹³, or OC(O)R¹³;

Each R¹¹ is independently C(O)R¹⁰, COOR¹⁰, or S(O)₂R¹⁰;

Each R¹² is independently H, C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, or phenyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF₃, OR¹³, SR¹³, NR¹³R¹³, COOR¹³, NO₂, CN, C(O)R¹³, C(O)NR¹³R¹³, NHC(O)R¹³, or OC(O)R¹³;

Each R¹³ is independently H; C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; C1-C10 alkyl optionally substituted with halo, CF₃, OR¹⁹, SR¹⁹, NR¹⁹R¹⁹, COOR¹⁹, NO₂, CN; or phenyl optionally substituted with halo, CF₃, OR¹⁹, SR¹⁹, NR¹⁹R¹⁹, COOR¹⁹, NO₂, CN;


Each R¹⁵ is independently C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R⁹; C1-C10 alkyl substituted with one or two independent aryl, R⁷ or R⁹ groups; or C1-C10 alkenyl substituted with aryl, R⁷ or R⁹;

Each R¹⁶ is independently C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; R⁹; C1-C10 alkyl substituted with one or two independent aryl, R⁷ or R⁹ groups; C1-C10 alkenyl substituted with aryl, R⁷ or R⁹; or phenyl substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, R⁹, halo, CF₃, OR¹², SR¹², NR¹²R¹², COOR¹², NO₂, CN, C(O)R¹², C(O)NR¹²R¹², NHC(O)R¹², NH(COOR¹²), S(O)₂NR¹²R¹², OC(O)R¹², C1-C10 alkyl substituted with R⁹, halo, CF₃, OR¹², SR¹², NR¹²R¹², COOR¹², NO₂, CN, C(O)R¹², C(O)NR¹²R¹², NHC(O)R¹², NH(COOR¹²), S(O)₂NR¹²R¹²;

Each haloalkyl is independently a C1-C10 alkyl substituted with one or more halogen atoms, selected from F, Cl, Br, or I, wherein the number of halogen atoms may not exceed that number that results in a perhaloalkyl group;

Each aryl is independently a 6-carbon monocyclic or 10-carbon bicyclic aromatic ring system optionally substituted with 1-3 independent C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; R^9 ; halo; haloalkyl; CF_3 ; OR^{12} ; SR^{12} ; $NR^{12}R^{12}$; $COOR^{12}$; NO_2 ; CN ; $C(O)R^{12}$; $C(O)C(O)R^{12}$; $C(O)NR^{12}R^{12}$; $S(O)_2R^{12}$; $N(R^{12})C(O)R^{12}$; $N(R^{12})(COOR^{12})$; $N(R^{12})S(O)_2R^{12}$; $S(O)_2NR^{12}R^{12}$; $OC(O)R^{12}$; $NR^{12}C(O)NR^{12}R^{12}$; $NR^{12}C(O)C(O)R^{12}$; $NR^{12}C(O)R^9$; $NR^{12}S(O)_2NR^{12}R^{12}$; $NR^{12}S(O)_2R^9$; $NR^{12}C(O)C(O)NR^{12}R^{12}$; C1-C10 alkyl substituted with 1-3 independent R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN , $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$; C2-C10 alkenyl substituted with 1-3 independent R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN , $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$; or R^{12} .

7. A compound of formula,



wherein,

R^2 is NR^5R^5 ; SR^5 ; OR^5 ; R^8 ; aryl; $N(R^5)-N=CH(R^8)$; $N(R^5)-N=CH(aryl)$; $NR^5-NR^5C(O)NR^5R^5$; $NR^5-NR^5R^{15}$; $NR^5-NR^5R^6$; C1-C10 alkyl substituted with aryl, R^8 , halo, CF_3 , SR^5 , OR^5 , $OC(O)R^5$, NR^5R^5 , NR^5R^6 , $COOR^5$, NO_2 , CN , $C(O)R^5$, $C(O)NR^5R^5$, or $S(O)_2NR^5R^5$; or C1-C10 alkenyl substituted with aryl, R^8 , halo, CF_3 , SR^5 , OR^5 , $OC(O)R^5$, NR^5R^5 , NR^5R^6 , $COOR^5$, NO_2 , CN , $C(O)R^5$, $C(O)NR^5R^5$, or $S(O)_2NR^5R^5$;

X is O or S;

R^4 is one, two, or three substituents, each independently selected from H, C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^8 ; halo; haloalkyl; CF_3 ; SR^5 ; OR^5 ; NR^5R^5 ; NR^5R^6 ; $COOR^5$; NO_2 ; CN; $C(O)R^5$; $C(O)C(O)R^5$; $C(O)NR^5R^5$; $OC(O)R^5$; $S(O)_2R^5$; $S(O)_2NR^5R^5$; $NR^5C(O)NR^5R^5$; $NR^5C(O)C(O)R^5$; $NR^5C(O)R^5$; $NR^5(COOR^5)$; $NR^5C(O)R^8$; $NR^5S(O)_2NR^5R^5$; $NR^5S(O)_2R^5$; $NR^5S(O)_2R^8$; $NR^5C(O)C(O)NR^5R^5$; $NR^5C(O)C(O)NR^5R^6$; C1-C10 alkyl substituted with aryl, R^7 or R^8 ; or C1-C10 alkenyl substituted with aryl, R^7 or R^8 ;

Each R^5 is independently H; C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^9 ; C1-C10 alkyl substituted with one or two independent aryl, R^7 or R^9 groups; C3-C10 cycloalkyl substituted with one or two independent aryl, R^7 or R^9 groups; or C1-C10 alkenyl substituted with aryl, R^7 or R^9 ;

Each R^6 is independently $C(O)R^5$, $COOR^5$, or $S(O)_2R^5$;

Each R^7 is independently halo, CF_3 , SR^{10} , OR^{10} , $OC(O)R^{10}$, $NR^{10}R^{10}$, $NR^{10}R^{11}$, $NR^{11}R^{11}$, $COOR^{10}$, NO_2 , CN, $C(O)R^{10}$, or $C(O)NR^{10}R^{10}$;

Each R^8 is independently a 5-8 membered monocyclic, 8-12 membered bicyclic, or 11-14 membered tricyclic ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if bicyclic, or 1-9 heteroatoms if tricyclic, said heteroatoms selected from O, N, or S, which may be saturated or unsaturated, and wherein 0, 1, 2 or 3 atoms of each ring may be substituted by a substituent independently selected from C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; aryl; R^9 ; halo; sulfur; oxygen; CF_3 ; haloalkyl; SR^5 ; OR^5 ; $OC(O)R^5$; NR^5R^5 ; NR^5R^6 ; NR^6R^6 ; $COOR^5$; NO_2 ; CN; $C(O)R^5$; $C(O)NR^5R^5$; C1-C10 alkyl substituted with R^7 , R^9 or aryl; C1-C10 alkenyl substituted with R^7 , R^9 or aryl;

Each R^9 is independently a 5-8 membered monocyclic, 8-12 membered bicyclic, or 11-14 membered tricyclic ring system comprising 1-3 heteroatoms if monocyclic, 1-6 heteroatoms if bicyclic, or 1-9 heteroatoms if tricyclic, said heteroatoms

C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; halo; sulfur; oxygen; CF₃; haloalkyl; SR¹⁰; OR¹⁰; NR¹⁰R¹⁰; NR¹⁰R¹¹; NR¹¹R¹¹; COOR¹⁰; NO₂; CN; C(O)R¹⁰; or C(O)NR¹⁰R¹⁰;

Each R¹⁰ is independently H, C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; haloalkyl; C1-C10 alkyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF₃, OR¹³, SR¹³, NR¹³R¹³, COOR¹³, NO₂, CN, C(O)R¹³, C(O)NR¹³R¹³, NHC(O)R¹³, or OC(O)R¹³; or phenyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF₃, OR¹³, SR¹³, NR¹³R¹³, COOR¹³, NO₂, CN, C(O)R¹³, C(O)NR¹³R¹³, NHC(O)R¹³, or OC(O)R¹³;

Each R¹¹ is independently C(O)R¹⁰, COOR¹⁰, or S(O)₂R¹⁰;

Each R¹² is independently H, C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, or phenyl optionally substituted with 1-3 independent C1-C10 alkyl, C2-C10 alkenyl, C2-C10 alkynyl, C3-C10 cycloalkyl, C4-C10 cycloalkenyl, halo, CF₃, OR¹³, SR¹³, NR¹³R¹³, COOR¹³, NO₂, CN, C(O)R¹³, C(O)NR¹³R¹³, NHC(O)R¹³, or OC(O)R¹³;

Each R¹³ is independently H; C1-C10 alkyl; C2-C10 alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; C1-C10 alkyl optionally substituted with halo, CF₃, OR¹⁹, SR¹⁹, NR¹⁹R¹⁹, COOR¹⁹, NO₂, CN; or phenyl optionally substituted with halo, CF₃, OR¹⁹, SR¹⁹, NR¹⁹R¹⁹, COOR¹⁹, NO₂, CN;

Each R¹⁸ is independently C1-C10 alkyl or both R¹⁸ may be taken together as a C2-C7 alkyl chain; wherein any R¹⁸ may optionally be substituted with 1-3 independent R⁷ or R⁸;

Each R¹⁹ is independently H; C1-C10 alkyl; C3-C10 cycloalkyl or phenyl;

Each haloalkyl is independently a C1-C10 alkyl substituted with one or more halogen atoms, selected from F, Cl, Br, or I, wherein the number of halogen atoms may not exceed that number that results in a perhaloalkyl group;

Each aryl is independently a 6-carbon monocyclic or 10-carbon bicyclic aromatic ring system optionally substituted with 1-3 independent C1-C10 alkyl; C2-C10

alkenyl; C2-C10 alkynyl; C3-C10 cycloalkyl; C4-C10 cycloalkenyl; R^9 ; halo; haloalkyl; CF_3 ; OR^{12} ; SR^{12} ; $NR^{12}R^{12}$; $COOR^{12}$; NO_2 ; CN ; $C(O)R^{12}$; $C(O)C(O)R^{12}$; $C(O)NR^{12}R^{12}$; $S(O)_2R^{12}$; $N(R^{12})C(O)R^{12}$; $N(R^{12})(COOR^{12})$; $N(R^{12})S(O)_2R^{12}$; $S(O)_2NR^{12}R^{12}$; $OC(O)R^{12}$; $NR^{12}C(O)NR^{12}R^{12}$; $NR^{12}C(O)C(O)R^{12}$; $NR^{12}C(O)R^9$; $NR^{12}S(O)_2NR^{12}R^{12}$; $NR^{12}S(O)_2R^9$; $NR^{12}C(O)C(O)NR^{12}R^{12}$; C1-C10 alkyl substituted with 1-3 independent R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN , $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$; C2-C10 alkenyl substituted with 1-3 independent R^9 , halo, CF_3 , OR^{12} , SR^{12} , $NR^{12}R^{12}$, $COOR^{12}$, NO_2 , CN , $C(O)R^{12}$, $C(O)NR^{12}R^{12}$, $NHC(O)R^{12}$, $NH(COOR^{12})$, $S(O)_2NR^{12}R^{12}$, $OC(O)R^{12}$; or R^{12} .

8. A composition comprising a compound according to any of claims 1-7 and a pharmaceutically acceptable carrier.

9. The composition according to claim 8, further comprising an additional therapeutic agent.

10. A method of treating a kinase mediated disease or disease symptoms in a mammal comprising administration to said mammal of a compound of claim 1.

11. The method of claim 10, wherein the mammal is a human.

12. A method of inhibiting kinase activity in a mammal comprising the step of administering to said mammal a compound of claim 1.


13. The method of claim 12, wherein said mammal is a human.

14. A method of treating disease or disease symptoms in a mammal comprising

16. A method of making a pharmaceutically useful composition comprising combining a compound of claim 1 with one or more pharmaceutically acceptable carriers.

17. The method of claim 16, further comprising combining an additional therapeutic agent.

18. A method of making a compound of claim 1 comprising reacting a pyrimidinone of the formula:



with an appropriate nucleophilic agent, wherein the groups in said formula are as defined in claim 1.

19. The method of claim 18, wherein R^1 is CN; R^3 is phenyl optionally substituted with 1-3 independent R^4 ; X is O; and R^5 is Me.